arXiv:1003.4742v1 [cond-mat.stat-mech] 24 Mar 2010

Anomalous Transport in Velocity Space, from Fokker-Planck to General Equation

S.A. Trigger

Joint Institute for High Temperatures,
Russian Academy of Sciences,
13/19, Izhorskaia Str.,
Moscow 125412, Russia;
Humboldt University,
Physics Institute, Newtonstr. 15,
D-12489 Berlin-Adlershof, Germany
email: satron@mail.ru

The problem of anomalous diffusion in momentum (velocity) space is considered based on the master equation and the appropriate probability transition function (PTF). The approach recently developed for coordinate space in [1], is applied with necessary modifications to velocity space. A new general equation for the time evolution of the momentum distribution function in momentum space is derived. This allows the solution of various problems of anomalous transport when the probability transition function (PTF) has a long tail in momentum space.

For the opposite cases of the PTF rapidly decreasing as a function of transfer momenta (when large transfer momenta are strongly suppressed), the developed approach allows us to consider strongly non-equilibrium cases of the system evolution. The stationary and non-stationary solutions are studied.

As an example, the particular case of the Boltzmann-type PT- function for collisions of heavy and light particles with the determined (prescribed) distribution function, which can be strongly non-equilibrium, is considered within the proposed general approach. The appropriate diffusion and friction coefficients are found. The Einstein relation between the friction and diffusion coefficients is shown to be violated in these cases.

PACS number(s): 52.27.Lw, 52.20.Hv, 05.40.Fb

I. INTRODUCTION

Interest in the anomalous diffusion is prompted by a large variety of applications, i.e., semiconductors, polymers, some granular systems, plasmas under specific conditions, various objects in biological systems, physical-chemical systems, and others.

Deviation from the linear in time dependence $< r^2(t) >\sim t$ of the mean-squared displacement has been experimentally observed, in particular, under essentially non-equilibrium conditions and in some disordered systems. The mean-squared separation of a pair of particles passively moving in a turbulent flow anomalously increases (according to the Richardson's law) with the third power of time [2]. For diffusion typical of glasses and related complex systems [3], the observed anomalous time dependence is slower than linear. These two types of anomalous diffusion are referred to as superdiffusion, $< r^2(t) >\sim t^\alpha$ $(\alpha > 1)$, and subdiffusion $(\alpha < 1)$, [4]. To describe these two diffusion regimes, a number of efficient models and methods have been proposed. The continuous time random walk (CTRW) model of Scher and Montroll [5], leading to strongly subdiffusive behavior, provides a basis for understanding photoconductivity in strongly disordered and glassy semiconductors. The Levy-flight model [6], leading to superdiffusion, describes various phenomena such as self-diffusion in micelle systems [7], reaction and transport in polymer systems [8], and has been even to the stochastic description of financial market indices [9]. For both cases, the so-called fractional differential equations in coordinate and time spaces were successfully applied [10].

However, recently a more general approach has been proposed in [1,11], which reproduces the results of the standard fractional differentiation method (when it is applicable) and allows the describe of more complicated cases of anomalous diffusion. In [12], this approach has also been applied to diffusion in a time-dependent external field.

In this paper, the problem of anomalous diffusion in momentum (velocity) space is considered. In spite of the formal similarity, diffusion in momentum space is very different physically from coordinate space diffusion. This is obvious because momentum conservation which takes place in momentum space has no analogy in coordinate space.

Some aspects of anomalous diffusion in velocity space have been considered in several papers [13-18]. In general, in comparison with anomalous diffusion in coordinate space, anomalous diffusion in velocity space is poorly studied. To our knowledge, there still is no corresponding way to describe anomalous diffusion in velocity space self-consistently .

In this paper, a new kinetic equation for anomalous diffusion in velocity space is derived (see also [19,20]) based on the appropriate expansion of the PTF (in spirit of the approach proposed in [1] for the diffusion in coordinate space) and some particular problems are studied on this basis.

The paper is organized as follows. The diffusion in velocity space for the cases of the normal and anomalous behavior of the PT-function is presented in Section II. Starting from argumentation based on the Boltzmann-type of the PTF, we derive a new kinetic equation which then can be applied not only to the Boltzmann-type processes, but also to a wide class of processes occurring in nature with other types of PT-functions. The particular cases of anomalous diffusion for hard spheres collisions with the specific power-type prescribed distribution function of light particles are analyzed in Section III. More general examples of anomalous diffusion are also considered. The non-stationary solution for the distribution function identical at $t = 0$ to the initial distribution is found. In Section IV, the Boltzmann-type equation is used to consider the effect light particle drift on the PT-function and on the kinetic equation structure. In Section V, a short review of anomalous diffusion in coordinate space is presented. The method and results of this paper can be easily extended to describe plasma-like systems and anomalous processes in energy space.

II. DIFFUSION IN VELOCITY SPACE ON THE BASIS OF THE MASTER-TYPE EQUATION

Let us now consider the main problem formulated in the introduction, namely, the description of diffusion in velocity space (V-space). Such a description would be based on a corresponding master equation describing the balance of grains coming at and from the point p at the instant t. The structure of this equation

$$\frac{df_g(\mathbf{p},t)}{dt} = \int d\mathbf{q}\left\{W(\mathbf{q},\mathbf{p}+\mathbf{q})f_g(\mathbf{p}+\mathbf{q},t) - W(\mathbf{q},\mathbf{p})f_g(\mathbf{p},t)\right\}. \tag{1}$$

is formally similar to the master equation Eq. (62) (see below) in coordinate space. Surely, for coordinate space, there is no conservation law similar to that in momentum space. The probability transition function $W(\mathbf{q},\mathbf{p}')$ (the transition rate) describes the probability that a grain with momentum $\mathbf{p}'$ (point $\mathbf{p}'$) passes from this point $\mathbf{p}'$ to the point $\mathbf{p}$ per unit time by transferring the momentum $\mathbf{q} = \mathbf{p}' - \mathbf{p}$ to the surrounding medium. Assuming initially that the characteristic transfer momentum $\mathbf{q}$ is smaller than $\mathbf{p}$, one may expand Eq. (1) in on $\mathbf{q}$ to the second order. Then, we arrive at the usual form of the Fokker-Planck equation for the density distribution function $f_g(\mathbf{p},t)$ (see, e.g., [21])

$$\frac{df_g(\mathbf{p},t)}{dt} = \frac{\partial}{\partial p_\alpha}\left[A_\alpha(\mathbf{p})f_g(\mathbf{p},t) + \frac{\partial}{\partial p_\beta}\left(B_{\alpha\beta}(\mathbf{p})f_g(\mathbf{p},t)\right)\right]. \tag{2}$$

$$A_\alpha(\mathbf{p}) = \int d^s q q_\alpha W(\mathbf{q},\mathbf{p}); \quad B_{\alpha\beta}(\mathbf{p}) = \frac{1}{2}\int d^s q q_\alpha q_\beta W(\mathbf{q},\mathbf{p}). \tag{3}$$

The coefficients A_α and $B_{\alpha\beta}$ describe the friction force and diffusion, respectively. The indices α and β correspond to the coordinate axes x, y, z (for dimension $s = 3$).

Since the velocity of heavy particles is low, the $\mathbf{p}$-dependence of the PTF can be neglected for calculating the diffusion term, which in this case is a constant $B_{\alpha\beta} = \delta_{\alpha\beta}B$. The constant B is equal to the integral

$$B = \frac{1}{2s}\int d^s q q^2 W(q), \tag{4}$$

where s is the dimension of velocity space (V-space0. Neglecting the $\mathbf{p}$-dependence of the PT-function, we arrive at the coefficient $A_\alpha = 0$ (while the diffusion coefficient is constant).

As is well known this neglect is wrong and the coefficient A_α for the Fokker-Planck equation can be determined using the assumption that argument that the stationary distribution function is Maxwellian. Then, we arrive at the standard relation between the coefficients $MTA_\alpha(p) = p_\alpha B$ (here M is the particle mass and T is temperature of the particles in the equilibrium state). This relation is an analogue of the Einstein relation in coordinate space. However, this argumentation is not applicable to systems far from equilibrium,.

To find the coefficients in the kinetic equation, which are appropriate for more general non-equilibrium situations, e.g., for slowly decreasing PT-functions or (and) for strongly non-equilibrium kernels W, let us use a more general way based on a certain small parameter (in the simplest case, e.g., on the difference of velocities of light and heavy particles for Boltzmann-type collisions). To calculate the function A_α we have take into account that the function $W(\mathbf{q},\mathbf{p})$ is a scalar and can depend only on variables $q, \mathbf{q}\cdot\mathbf{p}, p$. Expanding $W(\mathbf{q},\mathbf{p})$ in the product $\mathbf{q}\cdot\mathbf{p}$, one arrives at the approximate representation of the functions $W(\mathbf{q},\mathbf{p})$ and $W(\mathbf{q},\mathbf{p}+\mathbf{q})$,

$$W(\mathbf{q},\mathbf{p}) \simeq W(q) + \tilde{W}'(q)(\mathbf{q}\cdot\mathbf{p}) + \frac{1}{2}\tilde{W}''(q)(\mathbf{q}\cdot\mathbf{p})^2. \tag{5}$$

$$W(\mathbf{q},\mathbf{p}+\mathbf{q}) \simeq W(q) + \tilde{W}'(q)\,(\mathbf{q}\cdot\mathbf{p}) + \frac{1}{2}\tilde{W}''(q)(\mathbf{q}\cdot\mathbf{p})^2 + q^2\tilde{W}'(q), \tag{6}$$

Here we introduced the derivatives $\tilde{W}'(q) \equiv \partial W(q, \mathbf{q}\cdot\mathbf{p}, p)/\partial(\mathbf{qp}) \mid_{\mathbf{q}\cdot\mathbf{p}=0,\, p=0}$ and $\tilde{W}''(q) \equiv \partial^2 W(q, \mathbf{q}\cdot\mathbf{p}, p)/\partial(\mathbf{qp})^2 \mid_{\mathbf{q}\cdot\mathbf{p}=0,\, p=0}$.

Then, with necessary accuracy (which corresponds to the derivation of the well known Fokker-Planck equation) for the coefficient A_α we find

$$A_\alpha(\mathbf{p}) = \int d^s q q_\alpha q_\beta p_\beta \tilde{W}'(q) = p_\alpha \int d^s q q_\alpha q_\alpha \tilde{W}'(q) = \frac{p_\alpha}{s} \int d^s q q^2 \tilde{W}'(q) \tag{7}$$

If the equality $\tilde{W}'(q) = W(q)/2MT$ is fulfilled for the function $W(\mathbf{q},\mathbf{p})$, we arrive at the usual Einstein relation for the coefficients A_α and $B_{\alpha\beta}$

$$MTA_\alpha(\mathbf{p}) = p_\alpha B \tag{8}$$

Let us check this relation for Boltzmann collisions which are described by the PT-function $W(\mathbf{q},\mathbf{p}) = w_B(\mathbf{q},\mathbf{p})$ [11]:

$$w_B(\mathbf{q},\mathbf{p}) = \frac{2\pi}{\mu^2 q} \int_{q/2\mu}^{\infty} du\, u \frac{d\sigma}{do} \left[\arccos\left(1 - \frac{q^2}{2\mu^2 u^2}\right), u \right] f_b(u^2 + v^2 - \mathbf{q}\cdot\mathbf{v}/\mu), \tag{9}$$

where $(\mathbf{p} = M\mathbf{v})$, the quantities $d\sigma(\chi, u)/do$, μ and f_b are, respectively, the scattering differential cross-section, the mass and distribution function for light particles. The arguments χ and u in the cross-section $d\sigma/do$ are the angle of the particle scattering in the center-of- mass coordinate system and the velocity of light particles before collision. In Eq. (9), we took into account the approximate equalities for light and heavy particle scattering, $q^2 \equiv (\triangle\mathbf{p})^2 = p'^2(1 - cos\theta)$ and $\theta \simeq \chi$ (where $p' = \mu u$ is the light particle momentum before collision). For the equilibrium Maxwellian distribution f_b^0, the equality $\tilde{W}'(q) = W(q)/2MT$ is evident and we arrive at the ordinary Fokker-Planck equation in velocity space with the constant diffusion $D \equiv B/M^2$ and friction $\beta \equiv B/MT = DM/T$ coefficients which satisfy the Einstein relation.

The problem of the determination of the W-function and the respective Fokker-Planck equation for the situations close to equilibrium are discussed in detail in the review [22]. However, even for quasi-equilibrium regimes, where long tails of the PTF-functions are absent, the consideration in [22] is restricted to the hard sphere interaction. The non-equilibrium forms of the W-function and the drastic changes in the Fokker-Planck equation structure for these situations were not considered, as well as the specific situations, when the long tail of the W-function exists. Therefore, anomalous diffusion in [22] is absent. As is easy to see, in the case of the hard-sphere cross-section and the equilibrium Maxwellian distribution, f_b Eq. (9) leads to the same result for the W_B function as has been discussed in [22].

For some non-equilibrium (stationary or non-stationary) states, the PTF can have a long tail as a function of q. In this case, we have to derive a generalization of the Fokker-Planck equation in spirit of the consideration [1,11] for coordinate case. The necessity of this derivation arises since the diffusion and friction coefficients in the form Eqs. (4),(7) diverge in the limit of large q, if the kernels of the functions W, $\tilde{W}'$ exhibit the asymptotic behavior, $W(q) \sim 1/q^{\alpha}$ with $\alpha \leq s+2$ and (or) $\tilde{W}'(q) \sim 1/q^{\beta}$ with $\beta \leq s+2$.

Let us substitute the expansions for W (as an example, we choose $s=3$, the arbitrary s can be considered by the similar way) into Eq. (1). With an accuracy up to $(\mathbf{q}\cdot\mathbf{p})^2$, we find

$$\frac{df_g(\mathbf{p},t)}{dt} = \int d\mathbf{q}\{f_g(\mathbf{p}+\mathbf{q},t)[W(q) + \tilde{W}'(q)\,(\mathbf{q}\cdot\mathbf{p}) + \frac{1}{2}\tilde{W}''(q)(\mathbf{q}\cdot\mathbf{p})^2 + q^2\tilde{W}'(q)] - f_g(\mathbf{p},t)[W(q) + \tilde{W}'(q)\,(\mathbf{q}\cdot\mathbf{p}) + \frac{1}{2}\tilde{W}''(q)(\mathbf{q}\cdot\mathbf{p})^2]\} \tag{10}$$

After the Fourier-transformation $f_g(\mathbf{s},t) = \int \frac{d\mathbf{p}}{(2\pi)^3} exp(i\mathbf{pr}) f_g(\mathbf{p},t)$, Eq. (10) reads

$$\frac{df_g(\mathbf{s},t)}{dt} = A(s)f_g(\mathbf{s},t) + B_\alpha(s)\frac{\partial f_g(\mathbf{s},t)}{\partial \mathbf{s}_\alpha} + C_{\alpha\beta}(s)\frac{\partial^2 f_g(\mathbf{s},t)}{\partial \mathbf{s}_\alpha \partial \mathbf{s}_\beta} \tag{11}$$

It should be noted that the variable $\mathbf{s}_\alpha$, arisen in Eq. (11) has the dimension $\mathbf{p}_\alpha^{-1}$ and is a formal variable for the Fourier-transformation.

The coefficients in Eq. (11) are given by

$$A(\mathbf{s}) = \int d\mathbf{q}[exp(-i(\mathbf{qs})) - 1]W(q) = 4\pi \int_0^\infty dq q^2 \left[\frac{sin\,(qs)}{qs} - 1\right] W(q) \tag{12}$$

$$B_\alpha(\mathbf{s}) \equiv s_\alpha B(s);\; B(s) = -\frac{i}{s^2}\int d\mathbf{q}\mathbf{q}\mathbf{s}[exp(-i(\mathbf{qs})) - 1]\tilde{W}'(q) = \frac{4\pi}{s^2}\int_0^\infty dq q^2 \left[cos\,(qs) - \frac{sin(qs)}{qs}\right]\tilde{W}'(q) \tag{13}$$

$$C_{\alpha\beta}(s) \equiv s_\alpha s_\beta C(s) = -\frac{1}{2}\int d\mathbf{q} q_\alpha q_\beta [exp(-i(\mathbf{qs})) - 1]\tilde{W}''(q) \tag{14}$$

$$C(s) = -\frac{1}{2s^4}\int d\mathbf{q}(\mathbf{qs})^{\mathbf{2}}[exp(-i(\mathbf{qs})) - 1]\tilde{W}''(q) = \frac{2\pi}{s^2}\int_0^\infty dq q^4 \left[\frac{2sin(qs)}{q^3 s^3} - \frac{2cos\,(qs)}{q^2 s^2} - \frac{sin(qs)}{qs} + \frac{1}{3}\right]\tilde{W}''(q) \tag{15}$$

For the isotropic function $f(\mathbf{s}) = f(s)$, one can rewrite Eq. (11) as

$$\frac{df_g(s,t)}{dt} = A(s)f_g(s,t) + B(s)s\frac{\partial f_g(s,t)}{\partial s} + C(s)s^2\frac{\partial^2 f_g(s,t)}{\partial s^2} \tag{16}$$

In the case where the PT-function $W(q)$ and the functions $\tilde{W}'(q)$ and $\tilde{W}''(q)$ strongly decrease for large values of q, the exponents under the integrals in the functions $A(s)$, $B(s)$, and $C(s)$ can be expanded as

$$A(s) \simeq -\frac{s^2}{6}\int d\mathbf{q}\, q^2 W(q);\; B(s) \simeq -\frac{1}{3}\int d\mathbf{q}\, q^2 \tilde{W}'(q);\; C(s) \simeq 0. \tag{17}$$

Then, the simplified kinetic equation in velocity space PT-function (which is non-equilibrium in the general case) is written as

$$\frac{df_g(s,t)}{dt} = A_0 s^2 f_g(s) + B_0 s\frac{\partial f_g(s)}{\partial s}, \tag{18}$$

where $A_0 \equiv -1/6\int d\mathbf{q}\, q^2 W(q)$ and $B_0 \equiv -1/3\int d\mathbf{q}\, q^2 \tilde{W}'(q)$ is defined by Eq. (17).

For $C(s) = 0$ the stationary solution of Eq. (16) is given by

$$f_g(s) = Cexp\left[-\int_0^s ds'\frac{A(s')}{s'B(s')}\right] = Cexp\left[-\frac{A_0 s^2}{2B_0}\right] \tag{19}$$

The respective normalized stationary momentum distribution is written as

$$f_g(p) = \frac{N_g B_0^{3/2}}{(2\pi A_0)^{3/2}} exp\,[-\frac{B_0 p^2}{2A_0}] \tag{20}$$

Therefore, the constant in Eq. (19) is $C = N_g$, where N_g is the density of heavy particles. Equation (18) and the obtained distribution are the generalization (for non-equilibrium situations) of the standard Fokker-Planck equation to the case of normal diffusion in velocity space (see,e.g., [21]). The characteristic feature of these physical situations is the existence of the fixed (prescribed) kernel $W(\mathbf{q},\mathbf{p})$ which is defined, e.g., by a certain non-Maxwellian distribution of small particles f_b. To show this in other way, let us make the Fourier transformation of Eq. (11) with $C = 0$ and respective coefficients A and B_α,

$$\frac{df_g(\mathbf{p},t)}{dt} = -A_0\frac{\partial^2 f_g(\mathbf{p},t)}{\partial p^2} - B_0\frac{\partial(p_\alpha f_g(\mathbf{p},t))}{\partial p_\alpha}, \tag{21}$$

Therefore, we arrive at the Fokker-Planck-type equation with the friction coefficient $\beta \equiv -B_0$ and diffusion coefficient $D = -A_0/M^2$. In general, these coefficients (Eq. (17)) do not satisfy the Einstein relation.

In the case of equilibrium (e.g., $f_b = f_b^0$, see above) for the W-function the equality $\tilde{W}'(q) = W(q)/2MT_b$ is fulfilled. Then, with the necessary accuracy of the order μ/M, we find $A(s)/sB(s) \equiv A_0/B_0 = MT_b$. In this case, the Einstein relation between the diffusion and friction coefficients $D = \beta T/M$ exists, and the standard Fokker-Planck equation is valid.

In fact, the approximation $C(s) \simeq 0$ is practically always applicable due to the small parameter (e.g., μ/M for the Boltzmann-type PTF, see below, Sec.III). Therefore, the general kinetic equation (16) for the Fourier-transform of the velocity distribution function takes the form

$$\frac{df_g(\mathbf{s},t)}{dt} = A(\mathbf{s})f_g(\mathbf{s}) + B_\alpha(\mathbf{s})\frac{\partial f_g(\mathbf{s},t)}{\partial \mathbf{s}_\alpha}, \tag{22}$$

where the coefficients $A(\mathbf{s})$ and $B_\alpha(\mathbf{s})$ are determined by (12),(13), respectively.

III. THE MODELS OF ANOMALOUS DIFFUSION IN V - SPACE

Now we can calculate the coefficients for the models of anomalous diffusion. We will also estimate for the function W'' and the tensor $C_{\alpha,\beta}$; although, at the end we neglect these terms due to the small transfer momentum in the collision process.

At first, we calculate the simple model, i.e., the system of hard spheres with different masses m and $M \gg m$, $d\sigma/do = a^2/4$. Let us suppose that light particles in the model under consideration are described by the prescribed stationary distribution $f_b = n_b\phi_b/u_0^3$ (where n_b, ϕ_b, and u_0 are, respectively, the density, non-dimensional distribution, and characteristic velocity of light particles). The integration variable is defined by the equality $\xi \equiv (u^2 + v^2 - \mathbf{q}\cdot\mathbf{v}/\mu)/u_0^2$,

$$W_a(\mathbf{q},\mathbf{p}) = \frac{n_b a^2 \pi}{2\mu^2 u_0 q}\int_{(q^2/4\mu^2+v^2-\mathbf{q}\cdot\mathbf{v}/\mu)/u_0^2}^{\infty} d\xi \,\cdot \phi_b(\xi). \tag{23}$$

If the distribution $\phi_b(\xi) = 1/\xi^\gamma$ ($\gamma > 1$) has a long tail, we get

$$W_a(\mathbf{q},\mathbf{p}) = \frac{n_b a^2 \pi}{2\mu^2 u_0 q}\frac{\xi^{1-\gamma}}{(1-\gamma)}|_{\xi_0}^{\infty} = \frac{n_b a^2 \pi}{2\mu^2 u_0 q}\frac{\xi_0^{1-\gamma}}{(\gamma-1)}, \tag{24}$$

where $\xi_0 \equiv (q^2/4\mu^2 + v^2 - \mathbf{q}\cdot\mathbf{v}/\mu)/u_0^2$.

For the case $p = 0$, $\xi_0 \to \tilde{\xi}_0 \equiv q^2/4\mu^2 u_0^2$, and we arrive at the expression for the anomalous PT-function $W \equiv W_a$,

$$W_a(\mathbf{q},\mathbf{p}=\mathbf{0}) = \frac{n_b a^2 \pi}{2^{3-2\gamma}(\gamma-1)\mu^{4-2\gamma}u_0^{3-2\gamma}q^{2\gamma-1}} \equiv \frac{C_a}{q^{2\gamma-1}}. \tag{25}$$

To determine the transport process the structure and the kinetic equation in velocity space, one should also find the functions $\tilde{W}'(q)$ and $\tilde{W}''(q)$.

To find the functions $\tilde{W}'(\mathbf{q},\mathbf{p})$ and $\tilde{W}''(\mathbf{q},\mathbf{p})$ for $p \neq 0$, we use full-value $\xi_0 \equiv (q^2/4\mu^2 + p^2/M^2 - \mathbf{q}\cdot\mathbf{p}/M\mu)/u_0^2$ and the derivatives of these functions on $\mathbf{q}\cdot\mathbf{p}$ at $p = 0$, $\xi_0' = -1/M\mu u_0^2$, and $\xi_0'' = 0$. Then

$$\tilde{W}'(\mathbf{q},\mathbf{p}) \equiv \frac{n_b a^2 \pi}{2M\mu^3 u_0^3 q}\xi_0^{-\gamma}; \quad \tilde{W}''(\mathbf{q},\mathbf{p}) \equiv \frac{n_b a^2 \pi \gamma}{2M^2\mu^4 u_0^5 q}\xi_0^{-\gamma-1} \tag{26}$$

Therefore, for $p = 0$ ($\xi_0 \to \tilde{\xi}_0$) we obtain the functions

$$\tilde{W}'(q) \equiv \frac{(4\mu^2 u_0^2)^{\gamma} n_b a^2 \pi}{2M\mu^3 u_0^3 q^{2\gamma+1}}; \quad \tilde{W}''(q) \equiv \frac{(4\mu^2 u_0^2)^{\gamma+1} n_b a^2 \pi \gamma}{2M^2\mu^4 u_0^5 q^{2\gamma+3}} \tag{27}$$

The function $A(s)$, according to Eq. (12), is given by

$$A(s) \equiv 4\pi \int_0^{\infty} dq q^2 \left[\frac{sin\,(qs)}{qs} - 1\right] W(q) = 4\pi C_a \int_0^{\infty} dq \frac{1}{q^{2\gamma-3}} \left[\frac{sin(qs)}{qs} - 1\right] \tag{28}$$

Comparing the reduced equation (see below) in velocity space with diffusion in coordinate space (see the Appendix), we can establish the correspondence ($2\gamma - 1 \leftrightarrow \alpha$ and $W(q) = C/q^{2\gamma-1}$). This means that the convergence of the integral on the right side of Eq. (28) (3d case) is provided if $3 < 2\gamma - 1 < 5$ or $2 < \gamma < 3$. The inequality $\gamma < 3$ provides the convergence for small values of the variable q ($q \to 0$), and the inequality $\gamma > 2$ provides the convergence for $q \to \infty$.

We now establish the conditions of the convergence of the integrals for $B(s)$ and $C(s)$ as

$$B(s) = \frac{4\pi}{s^2} \int_0^{\infty} dq q^2 \left[cos\,(qs) - \frac{sin(qs)}{qs}\right] \tilde{W}'(q) \tag{29}$$

The convergence of the function $B(s)$ exists for small values of q if $\gamma < 2$ and for large values of q ($q \to \infty$ for $\gamma > 1/2$.

Finally, the convergence for the function $C(s)$ is defined by the equalities $\gamma < 2$ for small values of q and $\gamma > 1$ for large values of q

$$C(s) = \frac{2\pi}{s^2} \int_0^{\infty} dq q^4 \left[\frac{2sin(qs)}{q^3 s^3} - \frac{2cos\,(qs)}{q^2 s^2} - \frac{sin(qs)}{qs} + \frac{1}{3}\right] \tilde{W}''(q) \tag{30}$$

Therefore, in the case under consideration, the convergence of the functions A, B, and C for large large values of q is defined only by the convergence of the function A, which means $\gamma > 2$. To provide convergence for small values of q, it is sufficient to provide convergence for the functions $B(s)$ and C, which means $\gamma < 2$. Therefore, for the purely power behavior of the function $f_b(\xi)$, the simultaneous convergence of the coefficients A, B and C cannot be provided. However, in real physical models, the convergence of the coefficients in the integration region $q \to 0$ is always provided, e.g., by a finite value of v (see Eq. (24)) or due to the non-power behavior of the PT-function W for small values of q (compare with the examples of anomalous diffusion in coordinate space [1]). Therefore, in the model under consideration, the "anomalous diffusion in velocity space" for the power behavior of $W(q)$, $\tilde{W}'(q)$ and $\tilde{W}''(q)$ at large values of the variable q exists for the asymptotic behavior of the PT-function $W(q \to \infty) \sim 1/q^{2\gamma-1}$ if the inequality $\gamma > 2$ is fulfilled. In this case the integrals for the coefficients $A(s)$,$B(s)$ and $C(s)$ converge. At the same time, the expansion of the exponential function in Eqs. (12)-(15) under the integrals, which leads to the Fokker-Planck type kinetic equation, is invalid for the power-type kernels $W(\mathbf{q}, \mathbf{p})$.

Now let us consider the more general model for which we will not connect the functions $W(q)$, $\tilde{W}'(q)$, and $\tilde{W}''(q)$ with a concrete form of $W(\mathbf{q}, \mathbf{p})$ which is in general unknown. In this case one can suggest that the functions have the independent one from another power-type q-dependence.

As an example, this dependence can be taken as the power-type one for three functions $W(q) \equiv a/q^{\alpha}$, $\tilde{W}'(q) \equiv b/q^{\beta}$, and $\tilde{W}''(q) \equiv c/q^{\eta}$, where α, β and η are independent and positive. Then, as follows from the above consideration, the convergence of the function W exists if $5 > \alpha > 3$ (for asymptotically small and large q, respectively). For the function $\tilde{W}'(q)$, the convergence condition for asymptotically small and large values of q is provided if $5 > \beta > 2$, respectively. Finally, for the function $\tilde{W}''(q)$, the convergence condition is $7 > \eta > 5$ (for asymptotically small and large values of q, respectively).

For this example, the kinetic equation Eq. (11) reads

$$\frac{df_g(\mathbf{s},t)}{dt} = P_0 s^{\alpha-3} f(\mathbf{s},t) + s^{\beta-5} P_1 s_i \frac{\partial}{\partial s_i} f(\mathbf{s},t) + s^{\eta-7} P_2 s_i s_j \frac{\partial^2}{\partial s_i \partial s_j} f(\mathbf{s},t), \tag{31}$$

where

$$P_0 = 4\pi a \int_0^{\infty} d\zeta \zeta^{2-\alpha} \left[\frac{sin\,\zeta}{\zeta} - 1\right] \tag{32}$$

$$P_1 = 4\pi b \int_0^{\infty} d\zeta \zeta^{2-\beta} \left[cos\,\zeta - \frac{sin\zeta}{\zeta}\right] \tag{33}$$

$$P_2 = 4\pi c \int_0^\infty d\zeta \zeta^{4-\eta} \left[\frac{sin\zeta}{\zeta^3} - \frac{cos\,\zeta}{\zeta^2} - \frac{sin\zeta}{2\zeta} + \frac{1}{6} \right] \tag{34}$$

As is easy to show, $P_0 = -|P_0|\, sgn\, a < 0$ and $P_1 = -|P_1|\, sgn\, b$. Taking into account the isotropy in s-space we can rewrite Eq. (31) in the form

$$\frac{df_g(s,t)}{dt} = P_0 s^{\alpha-3} f(s,t) + s^{\beta-4} P_1 \frac{\partial}{\partial s} f(s,t) + s^{\eta-5} P_2 \frac{\partial^2}{\partial s^2} f(s,t), \tag{35}$$

Naturally, Eqs. (31),(35) can be formally rewritten in momentum (or in velocity) space in terms of fractional derivatives of various orders. Therefore, as is easy to see, for the purely power behavior of the functions $W(q)$, $\tilde{W}'(q)$ and $\tilde{W}''(q)$ the solution with the convergent coefficients exists for the values of the powers in the intervals mentioned above. In the case under consideration the universal type of the anomalous diffusion in velocity space exists if $5 > \alpha > 3$, $5 > \beta > 2$ and $7 > \eta > 5$. This takes place even in the cases, when the functions $W(q)$, $\tilde{W}'(q)$ and $\tilde{W}''(q)$ have not cut-off for small values of q. Surely, the general description is also valid for more complicated functions W, W', and W'', characterized by the non-power q-dependence for small values of q.

Now let us take into account an important circumstance; in general, there is a small parameter μ/M in the problem under consideration , which can simplify the description of velocity diffusion. As is easy to see, e.g., based on the particular cases (see Eq. (27)) for the convergent kernels of anomalous transport, the term with the second derivative $\tilde{W}''$ in general equations (11),(16) for the distribution $f_g(\mathbf{p})$ is small (in comparison with the term with the second derivative. The similar smallness is observed for the case of normal diffusion in velocity space. This smallness is of the order of the small ratio μ/M of particle masses.

Therefore, for the most physically important kernels describing anomalous velocity diffusion, the term with the second space derivative can be omitted and the non-stationary *general diffusion equation* is given by

$$\frac{df_g(\mathbf{s},t)}{dt} = A(s) f_g(\mathbf{s},t) + B_\alpha(s) \frac{\partial f_g(\mathbf{s},t)}{\partial \mathbf{s}_\alpha} \tag{36}$$

or, for the isotropic case,

$$\frac{df_g(s,t)}{dt} = A(s) f_g(s,t) + B(s) s \frac{\partial f_g(s,t)}{\partial s} \tag{37}$$

In the case of the purely power behavior, $W(q) = a/q^\alpha$ and $W'(q) = b/q^\beta$, we have, as above, $A(s) = P_0 s^{\alpha-3}$ and $sB(s) = P_1 s^{\beta-4}$ (with the inequalities $5 > \alpha > 3$ and $5 > \beta > 2$). The stationary solution of Eq. (37) (see, also (19)) for the case under consideration is written as

$$f_g^{St}(s) = Cexp\left[-\int^s ds'\frac{A(s')}{s'B(s')}\right] = Cexp\left[-\frac{P_0 s^{\alpha-\beta+2}}{P_1(\alpha-\beta+2)}\right], \tag{38}$$

where $5 > \alpha - \beta + 2 > 0$.

To find the solution in the isotropic non-stationary case, Eq. (37) should be written as

$$\frac{dX(s,t)}{dt} - B(s)s\frac{\partial}{\partial s}X(s,t) = A(s), \tag{39}$$

where $X(s,t) \equiv lnf_g(s,t)$. The general non-stationary solution of this equation can be written as the sum of the general solution of the homogeneous equation $Y(s,t)$ (Eq. (39), where the function $A(s)$ is taken zero),

$$Y(s,t) = \Phi(\xi),\ \ \xi \equiv t + \int_{s_0}^{s} ds'\frac{1}{s'B(s')}. \tag{40}$$

Here Φ is the arbitrary function of the variable ξ. The particular solution $Z(s,t)$ of the inhomogeneous equation Eq. (39) reads

$$Z(s,t) \equiv f_g^{St}(s) = -\int_{s_0}^{s} ds'\frac{A(s')}{s'B(s')} \tag{41}$$

Therefore, we find

$$f_g(s,t) = exp\left[X(s,t)\right] \equiv exp\left[Y + Z\right] = L(\xi)f_g^{St}(s), \tag{42}$$

where $L(\xi)$ is the arbitrary function of ξ, which should be found from the initial condition $f_g(s,t=0) \equiv \phi_0(s)$.

The variable $\xi \equiv \xi(s,t)$ equals

$$\xi(s,t) = t + \int_{s_0}^{s} ds'\frac{1}{s'B(s')} = t + \frac{s^{5-\beta}}{P_1(5-\beta)} + c, \tag{43}$$

where c is the arbitrary constant which can be omitted due to the presence of the arbitrary function L.

The general non-stationary solution for the case under consideration reads

$$f_g(s,t) = L\left(t + \frac{s^{5-\beta}}{P_1(5-\beta)}\right) exp\left[-\frac{P_0 s^{\alpha-\beta+2}}{P_1(\alpha-\beta+2)}\right]. \tag{44}$$

The inequalities for the combinations of the above coefficients are $5 > \alpha - \beta + 2 > 0$, $3 > 5 - \beta > 0$. The unknown function L can be found from Eq. (44) and the initial condition $f_g(s,0) \equiv \phi_g(s)$:

$$L\left(\frac{s^{5-\beta}}{P_1(5-\beta)}\right) exp\left[-\frac{P_0 s^{\alpha-\beta+2}}{P_1(\alpha-\beta+2)}\right] = \phi_g(s), \tag{45}$$

The function $\phi_g(s) = \int d^3 p \, exp(i\mathbf{ps}) \, f_g(p, t = 0)$ is the Fourier-component of the initial distribution in momentum space. Using the notation $\zeta \equiv s^{5-\beta}/[P_1(5-\beta)]$ (which means $s(\zeta) \equiv [P_1(5-\beta)\zeta]^{1/(5-\beta)}$), we find

$$L(\zeta) = \phi_g[s(\zeta)] \, exp \left\{ \frac{P_0 [s(\zeta)]^{\alpha-\beta+2}}{P_1(\alpha-\beta+2)} \right\}, \tag{46}$$

Therefore, the time-dependent solution is

$$f_g(s,t) = \phi_g[s(\zeta+t)] \, exp \left\{ \frac{P_0[s(\zeta+t)]^{\alpha-\beta+2}}{P_1(\alpha-\beta+2)} \right\} \, exp \left[-\frac{P_0 s^{\alpha-\beta+2}}{P_1(\alpha-\beta+2)} \right], \tag{47}$$

where we have to express the value $s(\zeta+t) \equiv [P_1(5-\beta)(\zeta+t)]^{1/(5-\beta)}$ as the evident function of variables s, t:

$$s(\zeta+t) \equiv [P_1(5-\beta)(\zeta+t)]^{1/(5-\beta)} \equiv [s^{5-\beta} + P_1(5-\beta)t]^{1/(5-\beta)}, \tag{48}$$

or, finally,

$$f_g(s,t) = \phi_g \left([s^{5-\beta} + P_1(5-\beta)\, t]^{1/(5-\beta)} \right) \, exp \left\{ \frac{P_0 \, [s^{5-\beta} + P_1 \, (5-\beta) \, t]^{(\alpha-\beta+2)/(5-\beta)} - P_0 \, s^{\alpha-\beta+2}}{P_1 \, (\alpha-\beta+2)} \right\} \tag{49}$$

It should be emphasized that the real solution for the fractional powers $1/(5-\beta)$ and (or) $(\alpha-\beta+2)/(5-\beta)$ exists only if $P_1 > 0$. The limit $t \to \infty$ for the solution can be identical to the stationary solution only for specific initial conditions. For these cases, the stationary solution can be non-equilibrium.

For the power dependence of the functions $W(q)$ and $\tilde{W}'(q)$, the equation (36) can be formally written in terms of fractional derivatives,

$$\frac{df_g(\mathbf{p},t)}{dt} = P_0 D^{\nu} f_g(\mathbf{p},t) - P_1(3+\gamma) D^{\gamma} f_g(\mathbf{p},t) + P_1 p_\alpha D_\alpha^{\gamma+1} f_g(\mathbf{p},t), \tag{50}$$

where $\nu \equiv \alpha - 3$, $\gamma \equiv \beta - 5$ and $D_\alpha^{\gamma+1} f_g(\mathbf{p},t) \equiv i \int \, d^3 \, s \, exp(-i\mathbf{ps})) s_\alpha s^\gamma f_g(\mathbf{s},t)$.

The particular case of the collision term for anomalous diffusion in velocity space, considered in [17], is derived in Appendix II on the basis of the general equations (36),(47).

IV. DIFFUSION MODEL BASED ON BOLTZMANN COLLISIONS WITH DRIFT

Let us consider the simplest case of the non-equilibrium but stationary distribution f_b, i.e., the shifted velocity distribution.

The evident generalization of the PT-function $w^d_B(\mathbf{q},\mathbf{p})$ for this case (characterizing by the drift velocity $\mathbf{u_d}$ in the distribution function f_b) is given by

$$w^d_B(\mathbf{q},\mathbf{p},\mathbf{u_d}) = \frac{2\pi}{\mu^2 q}\int_{q/2\mu}^{\infty} duu\cdot\frac{d\sigma}{do}\left[\arccos{(1-\frac{q^2}{2\mu^2u^2})}, u\right]\times$$
$$f_b(u^2+(\mathbf{v}-\mathbf{u_d})^2-\mathbf{q}\cdot(\mathbf{v}-\mathbf{u_d})/\mu), \tag{51}$$

where $\mathbf{p} = M\mathbf{v}$. Again, as in Section II, to find the coefficients in the kinetic equation, let us use the difference between the velocities of light and heavy particles. At the same time, the drift velocity u_d, generally speaking, is not small in comparison with the current characteristic velocities u and q/μ of small particles.

To calculate the function A_α, we have take into account that the scalar function $w^d_B(\mathbf{q},\mathbf{p},\mathbf{u_d})$ can in general be written in the form $W(\mathbf{q},\mathbf{p},\mathbf{u_d}) = W(q,p,u_d,l,\xi,\eta)$ (here $l\equiv(\mathbf{q}\cdot\mathbf{p_d}))$ and expanded on $\xi\equiv(\mathbf{q}\cdot\mathbf{p})$ and $\eta\equiv(M\mathbf{u_d}\cdot\mathbf{p})\equiv(\mathbf{p_d}\cdot\mathbf{p})$. In fact, it is the expansion in the velocity $\mathbf{v}=\mathbf{p}/M$, which is small in comparison with other characteristic velocities q/μ, u, and, in the general case, u_d. As shown above (for the case $u_d=0$), to arrive at the simple and solvable equation for the distribution function f_g, taking into account the smallness of v in comparison with characteristic velocities and v^2 in comparison with $(\mathbf{q}\cdot\mathbf{v}/\mu))$, we have approximate the function $W(q,p,u_d,l,\xi,\eta)\simeq W(q,u_d,l,\xi,\eta)$, because we are interested mainly in high values of q for anomalous transport. At the same time, after this approximation and expansion on ξ and η (for $u_d\neq 0$), we can arrive (for the special case of kernels with the purely power-type q-dependence for small q) at the divergence of some coefficients of the diffusion equation. A similar situation takes place also for anomalous diffusion in coordinate space. This divergence is absent for the realistic PT-functions, which provide the cut-off for small values of q. This cut-off has the physical nature and is not related to the approximation $W(\mathbf{q},\mathbf{p},\mathbf{u_d})\equiv W(q,p,u_d,l,\xi,\eta)\simeq W(q,v=0,u_d,l,\xi,\eta)W(q,u_d,l,\xi,\eta)$.

Let us expand the PT-function $W(q,p,u_d,l,\xi,\eta)$ in ξ and η similarly to the more simple case (5),

$$W(\mathbf{q},\mathbf{p},\mathbf{u_d}) = W(q,p,u_d,l,\xi,\eta)\simeq W_0(q,p,u_d,l)+\partial W/\partial\xi\mid_{\xi,\eta=0}\xi+\partial W/\partial\eta\mid_{\xi,\eta=0}\eta+$$
$$\frac{1}{2}\partial^2W/\partial\xi^2\mid_{\xi,\eta=0}\xi^2+\frac{1}{2}\partial^2W/\partial\eta^2\mid_{\xi,\eta=0}\eta^2+\partial^2W/\partial\xi\partial\eta\mid_{\xi,\eta=0}\xi\eta, \tag{52}$$

Here $W_0(q,p,u_d,l)\equiv W(q,p,u_d,l,\xi,\eta)\mid_{\xi,\eta=0}$. Then, introducing the functions $V_1(q,p,u_d,l)=\partial W/\partial\xi\mid_{\xi,\eta=0}, U_1(q,p,u_d,l)=\partial W/\partial\eta\mid_{\xi,\eta=0}, V_2(q,p,u_d,l)=\frac{1}{2}\partial^2W/\partial\xi^2\mid_{\xi,\eta=0}$

$, U_2(q,p,u_d,l) = \frac{1}{2}\partial^2 W/\partial\eta^2 \mid_{\xi,\eta=0}, W_2(q,p,u_d,l) = \partial^2 W/\partial\xi\partial\eta \mid_{\xi,\eta=0}$ we can rewrite Eq. (52) in the form

$$\begin{gathered} W(q,p,u_d,l,\xi,\eta) \simeq W_0(q,p,u_d,l) + V_1(\mathbf{q}\cdot\mathbf{p}) + \\ U_1(\mathbf{p_d}\cdot\mathbf{p}) + V_2(\mathbf{q}\cdot\mathbf{p})^2 + U_2(\mathbf{p_d}\cdot\mathbf{p})^2 + W_2(\mathbf{q}\cdot\mathbf{p})(\mathbf{p_d}\cdot\mathbf{p}) \simeq \\ W_0(q,p=0,u_d,l) + V_1(q,p=0,u_d,l)(\mathbf{q}\cdot\mathbf{p}) + U_1(q,p=0,u_d,l)(\mathbf{p_d}\cdot\mathbf{p}), \end{gathered} \tag{53}$$

Finally, we set $p = 0$ in the coefficients of Eq. (53) and omit the terms with the second order derivatives due to the existence of the small parameter (e.g., μ/M). This means only the terms of the order of $\sqrt{\mu/M}$ are essential in the expansion of $W(\mathbf{q},\mathbf{p},\mathbf{u_d})$. Let us calculate $W(\mathbf{q},\mathbf{p}+\mathbf{q},\mathbf{u_d})$ taking into account the difference between the values of the characteristic momenta $\mathbf{q},\mathbf{p}+\mathbf{q},\mathbf{u_d}$ or the characteristic values of velocities.

The appropriate expansion for the function $W(\mathbf{q},\mathbf{p}+\mathbf{q},\mathbf{u_d})$ with the necessary accuracy is written as

$$\begin{gathered} W(\mathbf{q},\mathbf{p}+\mathbf{q},\mathbf{u_d}) \simeq W(\mathbf{q},\mathbf{p},\mathbf{u_d}) + \{q_\alpha \partial/\partial p_\alpha + \frac{1}{2} q_\alpha q_\beta \frac{\partial^2}{\partial p_\alpha \partial p_\beta}\} W(\mathbf{q},\mathbf{p},\mathbf{u_d}) \simeq \\ W_0(q,p=0,u_d,l) + V_1(q,p=0,u_d,l)(\mathbf{q}\cdot\mathbf{p}) + U_1(q,p=0,u_d,l)(\mathbf{p_d}\cdot\mathbf{p}) + \\ V_1(q,p=0,u_d,l)\mathbf{q}^2 + U_1(q,p=0,u_d,l)(\mathbf{q}\cdot\mathbf{p_d}) \end{gathered} \tag{54}$$

or

$$\begin{gathered} W(\mathbf{q},\mathbf{p}+\mathbf{q},\mathbf{u_d}) \simeq W_0(q,p=0,u_d,l) + \\ V_1(q,p=0,u_d,l)[\mathbf{q}\cdot(\mathbf{p}+\mathbf{q})] + U_1(q,p=0,u_d,l)[\mathbf{p_d}\cdot(\mathbf{p}+\mathbf{q})]. \end{gathered} \tag{55}$$

Then the collision term of the kinetic equation can be given by the formula

$$\begin{aligned} \frac{df_g(\mathbf{p},t)}{dt} = \int d\mathbf{q}\{[&W_0(q,p=0,u_d,l) + V_1(q,p=0,u_d,l)(\mathbf{q}\cdot\mathbf{p}) + U_1(q,p=0,u_d,l)(\mathbf{p_d}\cdot\mathbf{p}) \\ &+V_1(q,p=0,u_d,l)\mathbf{q}^2 + U_1(q,p=0,u_d,l)(\mathbf{q}\cdot\mathbf{p_d})] \\ &f_g(\mathbf{p}+\mathbf{q},t) - [W_0(q,p=0,u_d,l) + V_1(q,p=0,u_d,l)(\mathbf{q}\cdot\mathbf{p}) + \\ &U_1(q,p=0,u_d,l)(\mathbf{p_d}\cdot\mathbf{p})]f_g(\mathbf{p},t)\} \end{aligned} \tag{56}$$

After the Fourier-transformation (10), the distribution function $f_g(\mathbf{s}) =$

$\int \frac{d\mathbf{p}}{(2\pi)^3} exp(i\mathbf{ps}) f_g(\mathbf{p}, t)$ can be written in the form

$$\frac{df_g(\mathbf{s},t)}{dt} = \int d\mathbf{q}\{exp(-i\mathbf{qs})[W_0(q,p=0,u_d,l) - iV_1(q,p=0,u_d,l)(\mathbf{q}\cdot\frac{\partial}{\partial\mathbf{s}}) - iU_1(q,p=0,u_d,l)(\mathbf{p_d}\cdot\frac{\partial}{\partial\mathbf{s}})]f_g(\mathbf{s},t) - [W_0(q,p=0,u_d,l) - iV_1(q,p=0,u_d,l)(\mathbf{q}\cdot\frac{\partial}{\partial\mathbf{s}}) - iU_1(q,p=0,u_d,l)(\mathbf{p_d}\cdot\frac{\partial}{\partial\mathbf{s}})]f_g(\mathbf{s},t)\} \tag{57}$$

Therefore, we arrive at the equation similar to Eq. (36), but with the evidently non-isotropic structure,

$$\frac{df_g(\mathbf{s},t)}{dt} = A_d(\mathbf{s},\mathbf{p_d})f_g(\mathbf{s},t) + \mathbf{B}_d(\mathbf{s},\mathbf{p_d})\frac{\partial}{\partial\mathbf{s}}f_g(\mathbf{s},t), \tag{58}$$

where the coefficients are expressed as the integrals

$$A_d(\mathbf{s},\mathbf{p_d}) = \int d\mathbf{q}[exp(-i\mathbf{qs}) - 1]W_0(q,p=0,u_d,l), \tag{59}$$

$$\mathbf{B}_d(\mathbf{s},\mathbf{p_d}) = -i\int d\mathbf{q}[exp(-i\mathbf{qs}) - 1]\{V_1(q,p=0,u_d,l)\mathbf{q} + U_1(q,p=0,u_d,l)\mathbf{p}_d\} \equiv \mathbf{s}B'_d(\mathbf{s},\mathbf{p_d}) + \mathbf{p_d}B''_d(\mathbf{s},\mathbf{p_d}). \tag{60}$$

This equation will be analyzed in detail in the separate paper.

V. CONCLUSIONS

In this paper, the problem of anomalous diffusion in momentum (velocity) space has been consistently considered. The new kinetic equation for anomalous diffusion in velocity space has been derived for the general case of the non-equilibrium probability transition function.

It should be emphasized that these types of anomalous diffusion are not related to the time dispersion (the non-local in time PT-function). Anomalous diffusion in velocity space, created by the non-local in time master equation has been considered in a few papers (see, e.g., [15],[16], [23],[24]) on the basis of the well-known Fokker-Planck kernel in p-space.

The similar problem with the non-local in time PT-functions, which at the same time have long q-tails, has been considered for diffusion in velocity space separately, based on the results obtained in this paper.

The evolution of the distribution function can lead to the stationary non-equilibrium limit or be asymptotically time-dependent. The model of anomalous diffusion in velocity space is described based on the respective expansion of the kernel in the master equation. The conditions of the convergence of the coefficients of the new kinetic equation have been found for particular cases. The wide variety of anomalous processes in velocity space exists, since two (C is usually small) different coefficients enter the general diffusion equation even in the isotropic case.

The examples of anomalous diffusion for heavy particles, based on the Boltzmann kernel with the prescribed equilibrium and non-equilibrium distribution function for the light particles have been studied. The PT-function for the Boltzmann kernel depends on the appropriate cross-section for scattering of two types of colliding particles and on the prescribed velocity distribution for one sort (light in the case under consideration) particles. The particular case of the hard sphere interaction has been considered. The Einstein relation is applicable only to the equilibrium PT-function (hence, the equilibrium prescribed distribution of light particles). In the general case, even for the stationary solution, the Einstein relation between the diffusion and friction coefficients is absent. For normal diffusion, the friction and diffusion coefficients have been explicitly found for the non-equilibrium case. The stationary and non-stationary (with an initial condition) solutions of the general kinetic equation have been found. In the equilibrium case, the known Fokker-Planck equation is reproduced as a particular case. We also have shortly reviewed anomalous diffusion in coordinate space.

The kinetic equation obtained in this paper is applicable to various systems in physics (including plasma physics [25]), chemistry, and biology.

VI. APPENDIX

Appendix I. Diffusion in the coordinate space on the basis of the master-type equation

Let us consider diffusion in coordinate space on the basis of the master equation which describes the balance of grains coming in and out the point r at the moment t. The structure of this equation (see, e.g., [1,11]) is formally similar to the master equation in momentum space, which has been derived above,

$$\frac{df_g(\mathbf{r},t)}{dt} = \int d\mathbf{r}' \left\{ W(\mathbf{r},\mathbf{r}')f_g(\mathbf{r}',\mathbf{t}) - W(\mathbf{r}',\mathbf{r})f_g(\mathbf{r},t) \right\}. \tag{61}$$

Surely, for coordinate space, there is no conservation law similar to that in momentum space. The probability transition $W(\mathbf{r},\mathbf{r}')$ describes the probability that a grain transfers from the point $\mathbf{r}'$ to the point $\mathbf{r}$ per unit time. We can rewrite this equation in the coordinates $\mathbf{u} = \mathbf{r}' - \mathbf{r}$ and $\mathbf{r}$ as

$$\frac{df_g(\mathbf{r},t)}{dt} = \int d\rho \left\{ W(\mathbf{u},\mathbf{r}+\mathbf{u})f_g(\mathbf{r}+\mathbf{u},t) - W(\mathbf{u},\mathbf{r})f_g(\mathbf{r},t) \right\}. \tag{62}$$

Assuming that characteristic displacements are small, one can expand Eq. (62) and arrive at the Fokker-Planck form of the equation for the density distribution $f_g(\mathbf{r},t)$,

$$\frac{df_g(\mathbf{r},t)}{dt} = \frac{\partial}{\partial r_\alpha}\left[A_\alpha(\mathbf{r})f_g(\mathbf{r},t) + \frac{\partial}{\partial r_\beta}\left(B_{\alpha\beta}(\mathbf{r})f_g(\mathbf{r},t) \right) \right]. \tag{63}$$

The coefficients A_α and $B_{\alpha\beta}$ describing the acting force and diffusion, respectively, can be written as the functionals of the PT-function in coordinate space W. For the dimension s, these coefficients have the form

$$A_\alpha(\mathbf{r}) = \int d^s\mathbf{u}\mathbf{u}_\alpha W(\mathbf{u},\mathbf{r}) \tag{64}$$

and

$$B_{\alpha\beta}(\mathbf{r}) = \frac{1}{2}\int d^s\mathbf{u}\mathbf{u}_\alpha\mathbf{u}_\beta W(\mathbf{u},\mathbf{r}). \tag{65}$$

For the isotropic case, the probability function depends on $\mathbf{r}$ and the magnitude u of the vector $\mathbf{u}$. For a homogeneous medium, when the r-dependence of the PT-function is absent, the coefficient $A_\alpha = 0$, while the diffusion coefficient is the constant $B_{\alpha\beta} = \delta_{\alpha\beta}B$. The constant B is the integral

$$B = \frac{1}{2s}\int d^s\mathbf{u}u^2W(u). \tag{66}$$

This consideration cannot be applied to the specific situations in which the integral in Eq. (66) is infinite. In this case we should examine the general transport equation (61). Let us now consider the problem for the homogeneous and isotropic case, when the PT-function depends only on u. As shown in [11], after the Fourier-transformation we arrive at the following representation of Eq. (61)

$$\frac{df_g(\mathbf{k},t)}{dt} = \int d^s\mathbf{u}\left[\exp(i\mathbf{k}\mathbf{u}) - 1\right]W(u)f_g(\mathbf{k},t) \equiv X(\mathbf{k})f_g(\mathbf{k},t), \tag{67}$$

where $X(\mathbf{k}) \equiv X(k)$. Let us assume the simple form of the PT-function with the power dependence on the distance $W(\rho) = C/u^{\alpha}$, where C is a constant and $\alpha > 0$. Such a singular dependence is typical for the jump diffusion probability in heteropolymers in solutions (see, e.g., [26], where different applications of anomalous diffusion are considered based on the fractional differentiation method). For the one-dimensional case (s=1), we find

$$X(k) \equiv -4 \int_0^{\infty} du\, sin^2\left(\frac{k\,u}{2}\right) W(u) = -2^{3-\alpha} C |k|^{\alpha-1} \int_0^{\infty} d\zeta \frac{sin^2 \zeta}{\zeta^{\alpha}}. \qquad (68)$$

For $1 < \alpha < 3$, this function is finite and written as

$$X(k) = -\frac{C\,\Gamma[(3-\alpha)/2]\,|k|^{\alpha-1}}{2^{\alpha}\sqrt{\pi}\,\Gamma(\alpha/2)(\alpha-1)}, \qquad (69)$$

where Γ is the Gamma-function. At the same time, the integral in Eq. (66) for such PT-functions is infinite, since normal diffusion is absent.

The procedure considered for the simplest cases of the power dependence of the PT-function is equivalent to the equation with fractional space differentiation [10,26],

$$\frac{df_g(x,t)}{dt} = C \Delta^{\mu/2} f_g(x,t), \qquad (70)$$

Here $\Delta^{\mu/2}$ is the fractional Laplacian, the linear operator whose action on the function $f(x)$ in Fourier space is described by the expression $\Delta^{\mu/2} f(x) = -(k^2)^{\mu/2} f(k) = -|k|^{\mu} f(k)$. In the case considered above, $\mu \equiv (\alpha - 1)$, where $0 < \mu < 2$. For more general PT-functions, which (for arbitrary values of the variable u) are not proportional to the power of u, the method described above is also applicable, although the fractional derivative method is not exist.

In the case of the purely power dependence of the PT-function, the non-stationary solution for the density distribution describes the so-called super-diffusion (or Levy flights). The solution of Eq. (70) in the Fourier space reads

$$f_g(k,t) = \exp(-C|k|^{\mu} t), \qquad (71)$$

which, in coordinate space, corresponds to the so-called symmetric Levy stable distribution

$$f_g(x,t) = \frac{1}{(kt)^{1/\mu}} L\left[\frac{x}{(kt)^{1/\mu}}; \mu, 0\right] \qquad (72)$$

(here we use the canonical notation [27]).

In the general case it follows from Eq. (67) that

$$f_g(k,t) = C_1 \exp[X(k)t], \tag{73}$$

with a certain constant C_1.

The consideration based on the PTF given in this Section allows us to avoid the fractional differentiation method and to consider more general physical situations of the non-power type of the probability transitions. Let us consider a simple example. Taking (for the one-dimensional case) the PT function $W(u)$ in the form

$$W(u) = C\frac{1-\exp[-\sigma u^p]}{u^\alpha}, \tag{74}$$

with $p > 0$, we arrive at the function $X(k)$:

$$X(k) = -2^{3-\alpha}C|k|^{\alpha-1}\int_0^\infty d\zeta \frac{\{1-\exp[-\sigma(2\zeta/|k|)^p\ ]\}\, sin^2\zeta}{\zeta^\alpha} \equiv -2^{3-\alpha}C|k|^{\alpha-1}T(\sigma/|k|^p,\alpha). \tag{75}$$

It is easy to see that the function $T(\sigma/|k|^p,\alpha)$ is finite for $1 < \alpha < p+3$, since for small distances at $p > 0$ divergence is also suppressed for the powers $\alpha > 3$. A simple calculation for $\alpha = 2$ and $p = 1$ leads to the following result which cannot be found by the usual fractional differentiation method,

$$T(\sigma/|k|,2) = \frac{\pi}{2} - \arctan(|k|/\sigma) + \frac{\sigma}{2|k|}\ln\left[1 + k^2/\sigma^2\right]. \tag{76}$$

The asymptotic behavior of the function $X(k)$ for $k \to 0$ (or $\sigma \to \infty$) is similar, as follows from Eq. (76), to the case $W(u) = C/u^\alpha$. For the case under consideration, $\alpha = 2$, the limit $X(k \to 0) \to -\pi Ck$. For large values of k ($k \to \infty$ or $\sigma \to 0$), we find $X(k) \to (\sigma/k)\ln(k/\sigma)$.

In the general case, the universal behavior of the function $X(k)$ is provided by the asymptotical properties of the PT-function for large distances. The necessary condition for divergence of the coefficient is $1 < \alpha < p+3$.

Appendix II. Anomalous velocity diffusion for the specific case B(s)=const, C(s)=0

Let us now consider formally the particular case of anomalous diffusion in velocity space, when the specific structure of PTF $W(\mathbf{q},\mathbf{p})$ provides a rapid (let say, exponential) decrease in the functions $\tilde{W}'(q)$ and $\tilde{W}''(q)$. Therefore, the exponential function under the integrals in the coefficients $B(s)$ and $C(s)$ can be expanded, which means $B(s) = B_0$ and $C(s) \simeq 0$

respectively. At the same time, the function $W(q) \equiv a/q^{\alpha}$ has a purely power dependence on q.

In this particular case, the kinetic equation Eq. (35) reads

$$\frac{df_g(\mathbf{s},t)}{dt} = P_0 s^{\alpha-3} f_g(\mathbf{s},t) + B_0 s_i \frac{\partial}{\partial s_i} f_g(\mathbf{s},t), \tag{77}$$

or, formally, in the momentum space

$$\frac{df_g(\mathbf{p},t)}{dt} = P_0 D^{\nu} f_g(\mathbf{p},t) - B_0 \frac{\partial}{\partial p_i}[p_i f_g(\mathbf{p},t)], \tag{78}$$

where $\nu \equiv (\alpha - 3)$ $(2 > \nu > 0)$, and we introduced the fractional differentiation operator $D^{\nu} f(\mathbf{p},t) \equiv \int d\mathbf{s} s^{\nu} exp(-i\mathbf{ps}) f(\mathbf{s},t)$ in momentum space. This is a very particular case of the general equations Eqs. (22),(35). In this case the result is similar to the particular phenomenological model [17]. The stationary solution of Eq. (77) is given by

$$f_g(s) = C exp\,[-\frac{P_0 s^{\nu}}{\nu B_0}] \tag{79}$$

$$f_g(p) = C \int d^3 s exp(-i\mathbf{ps}) exp\,[-\frac{P_0 s^{\nu}}{\nu B_0}] \equiv \frac{4\pi C}{p} \int_0^{\infty} ds s sin(ps) exp\,[-\frac{P_0 s^{\nu}}{\nu B_0}] \tag{80}$$

As an example, let us choose the case $\nu = 1$. Then, we find $f_g(\mathbf{p})$

$$f_g(p) = \frac{8\pi C' P_0}{B_0[(p^2 + 4P_0^2/B_0^2)]^2}. \tag{81}$$

In the case $\nu = 1$ the long tail of the distribution is proportional to p^{-4} and the distribution $f_g(p)$ corresponds with the Cauchy-Lorentz distribution. Normalization of the distribution $f_g(p)$ leads to the value $C = n_g/(2\pi)^3$, where $n_g = N_g/V$ is the average density of particles undergoing diffusion in velocity space. A similar approach can be taken for other types of anomalous diffusion in velocity space.

Acknowledgment

This paper has been reported as the invited talk at the Kiev International Conference devoted to the 100th anniversary of N.N. Bogolyubov, in September 2009.

I appreciate to the Organizing Committee and the Chairman A.G. Zagorodny for invitation, as well as, many participants for the useful discussions. I am grateful to E.A. Allahyarov, M. Bonitz, W. Ebeling, A.M. Ignatov, M.Yu. Romanovsky, and I.M. Sokolov for many stimulating discussions of anomalous diffusion problems. I am grateful to M.S. Karavaeva for her permanent supporting interest to this work.

This study was supported by the Netherlands Organization for Scientific Research (NWO) and the Russian Foundation for Basic Research, project no. 07-02-01464-a.

---

[1] S.A. Trigger, G.J.F. van Heijst and P.P.J.M. Schram, *Journal of Physics, Conference Series* **11**, 37 (2005).

[2] A.S. Monin and A.M. Yaglom, 1975 *Statistical Fluid Mechanics: Mechanics of Turbulence* Vol II (Cambridge: MIT Press)

[3] B. Rinn, P. Mass and J.-P. Bouchaud 2000 *Phys. Rev. Lett.* **84**, 5405 (2000)

[4] R. Metzler, and J. Klafter, *Phys. Rep.* **339**, 1 (2000).

[5] H. Scher and E.W. Montroll, *Phys. Rev.* **B12**, 2455 (1975)

[6] J. Klafter, M.F. Schlezinger and G. Zumofen, *Phys. Today* **49**, 33 (1996)

[7] A. Ott, J.-P. Bouchaud, D. Langevin and W. Urbach, *Phys. Rev. Lett.* **65**, 2201 (1994)

[8] I.M. Sokolov, J. May and A. Blumen, *Phys. Rev. Lett.* **79**, 857 (1997)

[9] P. Gopikrishnan, V. Plerou, L.A.N. Amaral, M. Meyer and H.E. Stanley, *Phys. Rev.* **E60**, 5305 (1999)

[10] B.J. West, M. Bologna, and P.Grigolini, *Physics of Fractal Operators*, (Springer-Verlag New York), 2003.

[11] S.A. Trigger, G.J.F. van Heijst and P.P.J.M. Schram, *Physica A* **347**, 77 (2005)

[12] S.A. Trigger, *Physics Letters A* **372**, 8, p. 1288 (2008)

[13] H.C. Fogedby, *Phys. Rev. Lett.* **73**, 2517 (1994)

[14] S. Jespersen, R. Metzler, H.C. Fogedby, *Phys. Rev.* **E 59**, 2736 (1999)

[15] E. Barkai, *Phys. Rev.* **E60**, 5305 (2003)

[16] E. Barkai, *Journal of statistical physics*, **115**, 1537 (2004)

[17] A. V. Chechkin, V. Yu. Gonchar, M. Szydlowski, *Phys. Plasmas* **9**, 78 (2002)

[18] W. Ebeling and M. Yu. Romanovsky, *Contr. Pl. Physics* 49, 479 (2009)

[19] S.A. Trigger, *Phys. Letters A* **374**, 134 (2009); ArXiv 0907.2793 v1, [cond-matt. stat.-mech.], 16 July 2009

[20] S.A. Trigger, Anomalous Kinetics in Velocity Space: equations and models, Arxiv 0909.1055

v1, [cond-matt. stat.-mech.], 5 September 2009.

[21] E.M. Lifshitz, L.P. Pitaevsky, Physical Kinetics, Nauka, Moscow (1979) [in Russian]

[22] M.R. Hoare, *Adv. Chem. Phys.* **20**, 135 (1971)

[23] H. Affan, R. Friedrich, S. Eule, *Phys. Rev.* E 80, 011137 (2009)

[24] S.A. Trigger. *Phys. Rev.* **E67**, 046403 (2003).

[25] S.A. Trigger, W. Ebeling, G.J.F. van Heijst, P.P.J.M. Schram, I.M. Sokolov, On anomalous diffusion in a plasma in velocity space, ArXiv 1002.2726 v1, [cond-matt. stat.-mech.], 13 February 2010

[26] D. Brockmann, and I.M. Sokolov, *Chemical Physics* **284**, 409 (2002)

[27] W. Feller, An Introduction to the Probability Theory and its Applications, vol. I and II, Willey, New York, 1971